

Comprehensive compilation of Randomized Controlled Trials (RCTs) involving Psychodynamic treatments (PDT)

Peter Lilliengren, PhD, assistant professor
Department of Psychology, Stockholm University, SWEDEN
<http://w3.psychology.su.se/staff/pe/i/indexeng.html>

Last updated: 2016-11-30

#	Reference (primary and secondary articles)	Patient sample	PDT-model	Comparison group(s)
1.	Abbass, A., Sheldon, A., Gyra, J., & Kalpin, A. (2008). Intensive short-term dynamic psychotherapy for DSM-IV personality disorders: a randomized controlled trial. <i>J Nerv Ment Dis</i> , 196, 211-216. http://www.ncbi.nlm.nih.gov/pubmed/18340256	Personality disorders (mixed)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990)	Minimal contact
2.	Ahola, P., Joensuu, M., Knekt, P., Lindfors, O., Saarinen, P., Tolmunen, T., ... Lehtonen, J. (2016). Effects of Scheduled Waiting for Psychotherapy in Patients With Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 0(0), 1. http://doi.org/10.1097/NMD.0000000000000616	Depression	Long-Term Psychodynamic Psychotherapy (At least 12 months, biweekly sessions, no reference to manual)	Waitlist (6 months)
3.	Ajlchi, B., Nejati, V., Town, J. M., Wilson, R., & Abbass, A. (2016). Effects of Intensive Short-Term Dynamic Psychotherapy on depressive symptoms and executive functioning in Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 204(7), 500–5. http://doi.org/10.1097/NMD.0000000000000518	Depression	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990)	Waitlist
4.	Alstrom, J. E., Nordlund, C. L., Persson, G., Harding, M., & Ljungqvist, C. (1984). Effects of four treatment methods on social phobic patients not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 97–110. http://www.ncbi.nlm.nih.gov/pubmed/6148843	Social anxiety disorder (SAD)	Supportive dynamic psychotherapy (Dewald, 1964)	Prolonged exposure (in vivo w. therapist)/ + relaxation therapy
5.	Alstrom, J. E., Nordlund, G., Persson, G., Harding, M., & Lundqvist, C. (1984). Effects of four treatment methods on agoraphobic women not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 1–17. http://www.ncbi.nlm.nih.gov/pubmed/6147066	Agoraphobia (in women not suitable for insight-oriented psychotherapy)	Supportive dynamic psychotherapy (Dewald, 1964)	Prolonged exposure (in vivo w. therapist)/ + relaxation therapy
6.	Amianto, F., Ferrero, A., Pierò, A., Cairo, E., Rocca, G., Simonelli, B., ... Fassino, S. (2011). Supervised team management, with or without structured psychotherapy, in heavy users of a mental health service with borderline personality disorder: a two-year follow-up preliminary randomized study. <i>BMC psychiatry</i> , 11(1), 181. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&tool=pmcentrez&rendertype=abstract	Borderline personality disorder	Sequential Brief Adlerian Psychodynamic Psychotherapy (SB-APP; Fassino, 2008; 2009)	Supervised team management

7.	<p>Andersson, G., Paxling, B., Roch-Norlund, P., Ostman, G., Norgren, A., Almlöv, J., ... Silverberg, F. (2012). Internet-Based Psychodynamic versus Cognitive Behavioral Guided Self-Help for Generalized Anxiety Disorder: A Randomized Controlled Trial. <i>Psychotherapy and psychosomatics</i>, 81(6), 344–355. http://www.ncbi.nlm.nih.gov/pubmed/22964540</p>	Generalized Anxiety Disorder (GAD)	IPDT (SUBGAP; Silverberg, 2005)	ICBT (Paxling, 2011)
8.	<p>Arnevik, E., Wilberg, T., Urnes, O., Johansen, M., Monsen, J. T., & Karterud, S. (2009). Psychotherapy for personality disorders: short-term day hospital psychotherapy versus outpatient individual therapy - a randomized controlled study. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 24(2), 71–8. http://www.ncbi.nlm.nih.gov/pubmed/19097870</p> <p>Arnevik, E., Wilberg, T., Urnes, Ø., Johansen, M., Monsen, J. T., & Karterud, S. (2010). Psychotherapy for Personality Disorders: 18 Months' Follow-Up of the Ullevål Personality Project. <i>Journal of Personality Disorders</i>, 24(2), 188–203. http://guilfordjournals.com/doi/pdf/10.1521/pedi.2010.24.2.188</p>	Personality disorders (mixed)	Short-Term Day Hospital Psychotherapy (DHP; no specific reference)	Outpatient Individual Psychotherapy
9.	<p>Bachar, E., Latzer, Y., Kreitler, S., & Berry, E. M. (1999). Empirical comparison of two psychological therapies. Self psychology and cognitive orientation in the treatment of anorexia and bulimia. <i>J Psychother Pract Res</i>, 8, 115-128. http://www.ncbi.nlm.nih.gov/pubmed/10079459</p>	Bulimia	Self Psychological Treatment (based on Kohut, 1984)	Cognitive Orientation Treatment (Kreitler & Kreitler, 1990)
10.	<p>Baldoni, F., Baldaro, B., & Trombini, G. (1995). Psychotherapeutic perspectives in urethral syndrome. <i>Stress Medicine</i>, 11(July 1994), 79–84. http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract</p>	Urethral syndrome	STDP (Malan, 1979; Davanloo, 1980)	TAU
11.	<p>Balottin, U., Ferri, M., Racca, M., Rossi, M., Rossi, G., Beghi, E., ... Termine, C. (2014). Psychotherapy versus usual care in pediatric migraine and tension-type headache: a single-blind controlled pilot study. <i>Italian Journal of Pediatrics</i>, 40(1), 6. http://www.ncbi.nlm.nih.gov/pubmed/24444140</p>	Idiopathic headache (children/adolescents 6-18 yrs)	STPP (Aliprandi et al, 1990)	TAU
12.	<p>Barber, J. P., Barrett, M. S., Gallop, R., Rynn, M. a, & Rickels, K. (2011). Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. <i>The Journal of clinical psychiatry</i>, 1–8. http://www.ncbi.nlm.nih.gov/pubmed/22152401</p> <p>Zilcha-Mano, S., Dinger, U., McCarthy, K. S., Barrett, M. S., & Barber, J. P. (2013). Changes in well-being and quality of life in a randomized trial comparing dynamic psychotherapy and pharmacotherapy for major depressive disorder. <i>Journal of affective</i></p>	Depression	Supportive Expressive Therapy (Luborsky, 1984)	1. Placebo 2. SSRI/SNRI

	<p>disorders, 1–5. http://www.ncbi.nlm.nih.gov/pubmed/24176534</p>			
13.	<p>Barkham, M., Rees, A., Shapiro, D. A., Stiles, W. B., Agnew, R. M., & Halstead, J., et al. (1996). Outcomes of time-limited psychotherapy in applied settings: replicating the Second Sheffield Psychotherapy Project. <i>J Consult Clin Psychol</i>, 64, 1079-1085. http://www.ncbi.nlm.nih.gov/pubmed/8916639</p>	Depression	Psychodynamic-interpersonal Therapy (Hobson, 1985)	CBT (no reference)
14.	<p>Barkham, M., Shapiro, D. A., Hardy, G. E., & Rees, A. (1999). Psychotherapy in two-plus-one sessions: outcomes of a randomized controlled trial of cognitive-behavioral and psychodynamic-interpersonal therapy for subsyndromal depression. <i>J Consult Clin Psychol</i>, 67, 201-211. http://www.ncbi.nlm.nih.gov/pubmed/10224730</p>	Depression (subsyndromal)	Psychodynamic-interpersonal Therapy (3 sessions; Barkham & Shapiro, based on Hobson, 1985)	CBT (Beck et al 1979; Beck, 1995)
15.	<p>Bassett, D. L., & Pilowsky, I. (1985). A study of brief psychotherapy for chronic pain. <i>J Psychosom Res</i>, 29, 259-264. http://www.sciencedirect.com/science/article/pii/0022399885900522</p>	Chronic pain	12 sessions PDT (no reference)	Supportive psychotherapy (no reference)
16.	<p>Bastos, A. G., Guimaraes, L. S. P., & Trentini, C. M. (2014). The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression. <i>Psychotherapy Research : Journal of the Society for Psychotherapy Research</i>, (July 2014), 1–13. http://www.ncbi.nlm.nih.gov/pubmed/25041333</p> <p>Bastos, A. G., Pinto Guimarães, L. S., & Trentini, C. M. (2013). Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. <i>Journal of Affective Disorders</i>, 1–10. http://www.ncbi.nlm.nih.gov/pubmed/24103853</p>	Depression	Long-term psychodynamic psychotherapy (LTTP; Gabbard, 2004; 2010)	<p>1. Fluoxetine alone</p> <p>2. LTTP + fluoxetine</p>
17.	<p>Bateman, A., & Fonagy, P. (1999). Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. <i>The American Journal of Psychiatry</i>, 156(10), 1563–9. http://www.ncbi.nlm.nih.gov/pubmed/10518167</p> <p>Bateman, A., & Fonagy, P. (2008). 8-Year Follow-Up of Patients Treated for Borderline Personality Disorder: Mentalization-Based Treatment Versus Treatment As Usual. <i>The American Journal of Psychiatry</i>, 165(5), 631–8. http://www.ncbi.nlm.nih.gov/pubmed/18347003</p>	Borderline personality disorder	Mentalization based treatment (MBT; Bateman & Fonagy, 2004)	TAU
18.	<p>Bateman, A., & Fonagy, P. (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. <i>Am J Psychiatry</i>, 166, 1355-1364. http://www.ncbi.nlm.nih.gov/pubmed/19833787</p>	<p>Borderline personality disorder</p> <p>Borderline and co-morbid Antisocial personality disorder (subsample in secondary)</p>	Mentalization based treatment (MBT; Bateman & Fonagy, 2004)	Structured Clinical Management (SCM)

	Bateman, A., O'Connell, J., Lorenzini, N., Gardner, T., & Fonagy, P. (2016). A randomised controlled trial of mentalization-based treatment versus structured clinical management for patients with comorbid borderline personality disorder and antisocial personality disorder. <i>BMC Psychiatry</i> , 16(1), 304. http://bmcpsychiatry.biomedcentral.com/articles/10.1186/s12888-016-1000-9	study)		
19.	Ben-Itzhak , S., Bluvstein, I., Schreiber, S., Aharonov-Zaig, I., Maor, M., Lipnik, R., & Bloch, M. (2012). The Effectiveness of Brief Versus Intermediate Duration Psychodynamic Psychotherapy in the Treatment of Adjustment Disorder. <i>Journal of Contemporary Psychotherapy</i> , 42(4), 249–256. http://www.springerlink.com/index/10.1007/s10879-012-9208-6	Adjustment disorder	Brief Psychodynamic Psychotherapy (12 sessions; Malan, 1979)	Brief Psychodynamic Psychotherapy (1 yrs; Malan, 1979)
20.	Beresnevaite , M. (2000) Exploring the benefits of group psychotherapy in reducing alexithymia in coronary heart disease patients: a preliminary study. <i>Psychotherapy and Psychosomatics</i> , 69, 117–122. http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract	Alexithymia (in patients with coronary heart disease)	Group psychotherapy (no reference)	Psychoeducation
21.	Beutel , M. E., Dippel, A., Szczepanski, M., Thiede, R., & Wiltink, J. (2006). Mid-Term Effectiveness of Behavioral and Psychodynamic Inpatient Treatments of Severe Obesity Based on a Randomized Study. <i>Psychotherapy and Psychosomatics</i> , 75(6), 337–345. http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract	Severe obesity	Individual in-patient PDT (Local program, no reference)	In-patient CBT (Local program, no reference)
22.	Beutel , M. E., Scheurich, V., Knebel, A., Michal, M., Wiltink, J., Graf-Morgenstern, M., ... Subic-Wrana, C. (2013). Implementing panic-focused psychodynamic psychotherapy into clinical practice. <i>Canadian journal of psychiatry. Revue canadienne de psychiatrie</i> , 58(6), 326–34. http://www.ncbi.nlm.nih.gov/pubmed/23768260	Panic disorder	Panic Focused Psychodynamic Psychotherapy (PFPP; Milrod)	CBT
23.	Beutel , M. E., Weissflog, G., Leuteritz, K., Wiltink, J., Haselbacher, a., Ruckes, C., ... Brahler, E. (2013). Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: results of a randomized controlled multicenter trial. <i>Annals of Oncology</i> , 1–7. http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526 Weißflog , G., Brähler, E., Leuteritz, K., Barthel, Y., Kuhnt, S., Wiltink, J., ... Beutel , M. E. (2015). Does psychodynamic short-term psychotherapy for depressed breast cancer patients also improve fatigue? Results from a randomized controlled trial. <i>Breast Cancer Research and Treatment</i> . http://link.springer.com/10.1007/s10549-015-3494-0	Depression (in breast cancer patients)	Supportive-expressive therapy (Haselbacher, Barthel, Brähler et al., 2010; based on Luborsky, 1984)	TAU
24.	Bloch , M., Meiboom, H., Lorberblatt, M., Bluvstein, I., Aharonov, I., & Schreiber, S.	Depression (post-partum)	Brief Dynamic Psychotherapy (BDT;	Brief Dynamic Psychotherapy (BDT;

	(2012). The effect of sertraline add-on to brief dynamic psychotherapy for the treatment of postpartum depression: a randomized, double-blind, placebo-controlled study. <i>The Journal of clinical psychiatry</i> , 73(2), 235–41. http://www.ncbi.nlm.nih.gov/pubmed/22401479		Malan, 1979) + placebo meds	Malan, 1979) + Sertraline
25.	Bögels, S., Wijts, P., Oort, F., & Sallaerts, S. J. (2014). Psychodynamic psychotherapy versus cognitive-behavior therapy for social anxiety disorder: An efficacy and partial effectiveness trial. <i>Depression and Anxiety</i> , 31(5), 363–373. http://doi.wiley.com/10.1002/da.22246	Social anxiety disorder (SAD)	STDP (Malan, 1979; also Luborsky 1984)	CBT (Clark & Wells, 1995)
26.	Bressi, C., Porcellana, M., Marinaccio, P. M., Nocito, E. P., & Magri, L. (2010). Short-term psychodynamic psychotherapy versus treatment as usual for depressive and anxiety disorders: a randomized clinical trial of efficacy. <i>The Journal of Nervous and Mental Disease</i> , 198(9), 647–52. http://www.ncbi.nlm.nih.gov/pubmed/20823726	Mixed sample (depression and anxiety)	Brief Dynamic Psychotherapy (BDT; Malan, 1976; Osimo & Malan, 1992)	TAU
27.	Brockman, B., Poynton, A., Ryle, A., & Watson, J. P. (1987). Effectiveness of time-limited therapy carried out by trainees. Comparison of two methods. <i>Br J Psychiatry</i> , 151, 602-610. http://bjp.rcpsych.org/content/151/5/602	Mixed sample	Interpretive therapy (Mann & Goldman, 1982)	Cognitive-analytic therapy (CAT; Ryle, 1979; 1982)
28.	Brodady, H., & Andrews, G. (1983). Brief psychotherapy in family practice. A controlled prospective intervention trial. <i>Br J Psychiatry</i> , 143, 11-19. http://bjp.rcpsych.org/content/143/1/11.short	Mixed sample	STDP (Balant, Malan & Sifneos)	1. Family practitioner 2. No treatment
29.	Brom, D., Kleber, R. J., & Defares, P. B. (1989). Brief psychotherapy for posttraumatic stress disorders. <i>J Consult Clin Psychol</i> , 57, 607-612. http://www.ncbi.nlm.nih.gov/pubmed/2571625	PTSD	Brief Psychodynamic Therapy (Horowitz, 1976)	1. Trauma Desensitization (no specific reference) 2. Hypnotherapy (no specific reference)
30.	Budman, S. H., Demby, A., Redondo, J. P., Hannan, M., Feldstein, M., & Ring, J., et al. (1988). Comparative outcome in time-limited individual and group psychotherapy. <i>Int J Group Psychother</i> , 38, 63-86. http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085	Mixed sample	Time-limited individual psychotherapy (Mann, 1973)	Time-limited group psychotherapy (Budman & Bennett, 1983)
31.	Burnand, Y., Andreoli, A., Kolatte, E., Venturini, A., & Rosset, N. (2002). Psychodynamic psychotherapy and clomipramine in the treatment of major depression. <i>Psychiatr Serv</i> , 53, 585-590. http://www.ncbi.nlm.nih.gov/pubmed/11986508	Depression	PDT + clomipramine , (unclear which model, Safran & Muran is mentioned among references)	Supportive care + clomipramine
32.	Burns, A., Guthrie, E., Marino-Francis, F., Busby, C., Morris, J., Russell, E., ... Byrne, J. (2005). Brief psychotherapy in Alzheimer's disease. <i>British Journal Of Psychiatry</i> , 187(2), 143–147. http://bjp.rcpsych.org/content/187/2/143.short	Alzheimers disease	Psychodynamic-Interpersonal therapy (Hobson, 1985)	TAU
33.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). A Randomized Double-Blind Controlled Trial Comparing Davanloo's	Medically unexplained pain (MUP)	Internet-delivered ISTDP (Abbass, 2015)	TAU

	Intensive Short-Term Dynamic Psychotherapy as Internet-Delivered Versus Treatment as Usual for Medically Unexplained Pain: A Six-Month Pilot Study. <i>Psychosomatics</i> , 57(3), 292–300. http://linkinghub.elsevier.com/retrieve/pii/S0033318216000025			
34.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). Efficacy of Intensive Short-Term Dynamic Psychotherapy for Medically Unexplained Pain: A Pilot Three-Armed Randomized Controlled Trial Comparison with Mindfulness-Based Stress Reduction. <i>Psychotherapy and Psychosomatics</i> , 123–125. http://www.karger.com/?doi=10.1159/000441698	Medically unexplained pain (MUP)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	1. Mindfulness-based stress reduction (MBSR; no reference) 2. TAU
35.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). Telemedicine vs. in-person delivery of intensive short-term dynamic psychotherapy for patients with medically unexplained pain: A 12-month randomized, controlled trial. <i>Journal of Telemedicine and Telecare</i> , 0(0), 1–9. http://itt.sagepub.com/lookup/doi/10.1177/1357633X15627382	Medically unexplained pain (MUP)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	Internet-delivered ISTDP (ID-ISTDP)
36.	Clarici, A., Pellizzoni, S., Guaschino, S., Alberico, S., Bembich, S., Giuliani, R., ... Panksepp, J. (2015). Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study. <i>Frontiers in Psychology</i> , 06, 1–10. http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract	Depression (post-partum)	Psychodynamic approach (non-manualized?) + oxytocine	Psychodynamic approach (non-manualized?) + placebo
37.	Clarkin, J. F., Levy, K. N., Lenzenweger, M. F., & Kernberg, O. F. (2007). Evaluating three treatments for borderline personality disorder: a multiwave study. <i>Am J Psychiatry</i> , 164, 922-928. http://www.ncbi.nlm.nih.gov/pubmed/17541052 Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., & Kernberg, O. F. (2006). Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. <i>Journal of Consulting and Clinical Psychology</i> , 74(6), 1027–40. http://www.ncbi.nlm.nih.gov/pubmed/17154733	Borderline personality disorder	Transference-Focused Psychotherapy (TFP; Kernberg)	1. DBT (Linehan, 1991) 2. Supportive treatment (Rockland, 1992)
38.	Connolly Gibbons, M. B., Thompson, S. M., Scott, K., Schauble, L. a, Mooney, T., Thompson, D., ... Crits-Christoph, P. (2012). Supportive-expressive dynamic psychotherapy in the community mental health system: A pilot effectiveness trial for the treatment of depression. <i>Psychotherapy (Chicago, Ill.)</i> , 49(3), 303–16. http://www.ncbi.nlm.nih.gov/pubmed/22962971	Depression	Supportive-expressive therapy (manual adapted from Luborsky, 1984)	TAU
39.	Connolly Gibbons, M. B., Gallop, R.,	Depression	Supportive-expressive	CBT (Beck, 1995)

	Thompson, D., Luther, D., Crits-Christoph, K., Jacobs, J., ... Crits-Christoph, P. (2016). Comparative effectiveness of Cognitive Therapy and Dynamic Psychotherapy for Major Depressive Disorder in a community mental health setting. <i>JAMA Psychiatry</i> , 19104, 1–8. http://doi.org/10.1001/jamapsychiatry.2016.1720		therapy (Book, 1998; Luborsky, 1984)	
40.	Cooper, P. J., Murray, L., Wilson, A., & Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. <i>Br J Psychiatry</i> , 182, 412-419. http://www.ncbi.nlm.nih.gov/pubmed/12724244 Murray, L., Cooper, P. J., Wilson, A., & Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post- partum depression: 2. Impact on the mother-child relationship and child outcome. <i>British Journal of Psychiatry</i> , 182, 420–7. http://www.ncbi.nlm.nih.gov/pubmed/12724245	Depression (post-partum)	Psychodynamic therapy (Cramer et al, 1990; Stern, 1995)	1. TAU 2. Non-directive counseling (Holden et al, 1989) 3. Cognitive-Behavioural Therapy (Hawton et al, 1989)
41.	Covi, L., & Lipman, R. S. (1987). Cognitive behavioral group psychotherapy combined with imipramine in major depression. <i>Psychopharmacol Bull</i> , 23, 173-176. http://www.ncbi.nlm.nih.gov/pubmed/3602315	Depression	Traditional Group Therapy (based on “psychodynamic-Interpersonal” theories but with no reference)	1. CBT group therapy (Beck, 1979; Covi, 1982) 2. CBT group therapy + Imipramine
42.	Covi, L., Lipman, R. S., Derogatis, L. R., Smith, J. E., & Pattison, J. H. (1974). Drugs and group psychotherapy in neurotic depression. <i>Am J Psychiatry</i> , 131, 191-198. http://ajp.psychiatryonline.org/doi/abs/10.1176/ajp.131.2.191	Depression (chronic)	Psychodynamically oriented group therapy (no reference) + medication or placebo	Brief supportive contact + medication or placebo
43.	Creed, F., Fernandes, L., Guthrie, E., Palmer, S., Ratcliffe, J., & Read, N., et al. (2003). The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. <i>Gastroenterology</i> , 124, 303-317. http://www.ncbi.nlm.nih.gov/pubmed/12557136 Hyphantis, T., Guthrie, E., Tomenson, B., & Creed, F. (2009). Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome. <i>Pain</i> , 145(1-2), 196–203. http://www.ncbi.nlm.nih.gov/pubmed/19643544	Irritable bowel syndrome	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU/SSRI
44.	Crits-Christoph, P., Gibbons, M. B. C., Narducci, J., Schamberger, M., & Gallop, R. (2005). Interpersonal Problems and the Outcome of Interpersonally Oriented Psychodynamic Treatment of GAD. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 42(2), 211–224. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211	Generalized anxiety disorder (GAD)	Supportive Expressive Therapy (Luborsky (1984)	Supportive therapy (Borkovec & Matthews, 1988; Borkovec & Costello, 1993)
45.	Crits-Christoph, P., Siqueland, L., Blaine, J., Frank, A., Luborsky, L., & Onken, L. S., et al.	Cocaine dependence	Supportive Expressive Therapy (Luborsky (1984)	1. Cognitive therapy (Beck, 1993) + group drug

	<p>(1999). Psychosocial treatments for cocaine dependence: National Institute on Drug Abuse Collaborative Cocaine Treatment Study. <i>Arch Gen Psychiatry</i>, 56, 493-502. http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777</p> <p>Crits-Christoph, P., Gibbons, M. B. C., Gallop, R., Ring-Kurtz, S., Barber, J. P., Worley, M., ... Hearon, B. (2008). Supportive-Expressive Psychodynamic Therapy for Cocaine Dependence: A Closer Look. <i>Psychoanalytic Psychology: The Official Journal of the Division of Psychoanalysis, American Psychological Association, Division 39</i>, 25(3), 483-498. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&tool=pmcentrez&rendertype=abstract</p>		+ group drug counseling (Mercer & Woody, 1992)	counseling (Mercer & Woody, 1992) 2. Group drug counseling alone
46.	<p>Cummings, J. G., & Wittenberg, J. V. (2008). Supportive expressive therapy - Parent child version: an exploratory study. <i>Psychotherapy Theory, Research, Practice, Training</i>, 45, 148-164. http://www.ncbi.nlm.nih.gov/pubmed/22122414</p>	Oppositional defiant disorder/disruptive behavior disorder (children)	Supportive Expressive Therapy – Parent Child (SET-PC; Wittenberg, adapted from Luborsky, 19854)	The Incredible Years Parenting Program
47.	<p>Dare, C., Eisler, I., Russell, G., Treasure, J., & Dodge, L. (2001). Psychological therapies for adults with anorexia nervosa: randomised controlled trial of out-patient treatments. <i>Br J Psychiatry</i>, 178, 216-221. http://www.ncbi.nlm.nih.gov/pubmed/11230031</p>	Anorexia	Focal psychoanalytic psychotherapy (Dare, 1995; Malan, 1979)	Cognitive Analytic Therapy (Ryle, 1990; Treasure et al, 1995) Family therapy (no specific reference)
48.	<p>de Jonghe, F., Hendriksen, M., van Aalst, G., Kool, S., Peen, J., & Van, R., et al. (2004). Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. <i>British Journal of Psychiatry</i>, 185, 37-45. http://www.ncbi.nlm.nih.gov/pubmed/15231554</p>	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005)	Short Psychodynamic Supportive Psychotherapy + antidepressants
49.	<p>de Jonghe, F., Kool, S., van Aalst, G., Dekker, J., & Peen, J. (2001). Combining psychotherapy and antidepressants in the treatment of depression. <i>Journal of Affective Disorders</i>, 64(2-3), 217-29. http://www.ncbi.nlm.nih.gov/pubmed/11313088</p>	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) + medication	Medication alone
50.	<p>de Roten, Y., Ambresin, G., Herrera, F., Fassassi, S., Fournier, N., Preisig, M., & Despland, J.-N. (2017). Efficacy of an adjunctive brief psychodynamic psychotherapy to usual inpatient treatment of depression: Results of a randomized controlled trial. <i>Journal of Affective Disorders</i>, 209, 105-113. http://doi.org/10.1016/j.jad.2016.11.013</p>	Depression	Intensive Brief Psychodynamic Psychotherapy (IBPP; Despland et al., 2010) + TAU	TAU only
51.	<p>Dekker, J. J. M., Koelen, J. A., Van, H. L., Schoevers, R. A., Peen, J., Hendriksen, M., ... De Jonghe, F. (2008). Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. <i>Journal of Affective Disorders</i>, 109(1-2), 183-188. http://linkinghub.elsevier.com/retrieve/pii/S0</p>	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005)	Pharmacotherapy

	165032707003588 Dekker, J., Van, H. L., Hendriksen, M., Koelen, J., Schoevers, R. A., Kool, S., ... Peen, J. (2013). What is the best sequential treatment strategy in the treatment of depression? Adding pharmacotherapy to psychotherapy or vice versa? <i>Psychotherapy and Psychosomatics</i> , 82(2), 89–98. http://www.karger.com/doi/10.1159/000341177			
52.	Dekker, J., Molenaar, P. J., Kool, S., Van Aalst, G., Peen, J., & de Jonghe, F. (2005). Dose-effect relations in time-limited combined psycho-pharmacological treatment for depression. <i>Psychological Medicine</i> , 35(1), 47–58. http://www.journals.cambridge.org/abstract/S0033291704002685	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 8 sessions + medication	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 16 sessions + medication
53.	Doering, S., Hörz, S., Rentrop, M., Fischer-Kern, M., Schuster, P., & Benecke, C., et al. (2010). Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. <i>Br J Psychiatry</i> , 196, 389-395. http://www.ncbi.nlm.nih.gov/pubmed/20435966	Borderline personality disorder	Transference-focused psychotherapy (TFP; Kernberg)	TAU (experienced therapists)
54.	Driessen, E., Van, L. H., Don, F. J., Peen, J., Kool, S., Westra, D., ... Dekker, J. J. M. (2013). The Efficacy of Cognitive-Behavioral Therapy and Psychodynamic Therapy in the Outpatient Treatment of Major Depression: A Randomized Clinical Trial. <i>American Journal of Psychiatry</i> , 170(9), 1041-1050 http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899 Driessen, E., Van, H. L., Peen, J., Don, F. J., Kool, S., Westra, D., ... Dekker, J. J. (2014). Therapist-rated outcomes in a randomized clinical trial comparing cognitive behavioral therapy and psychodynamic therapy for major depression. <i>Journal of Affective Disorders</i> , 170C, 112–118. http://www.ncbi.nlm.nih.gov/pubmed/25240140	Depression	Short-term psychodynamic supportive psychotherapy (de Jonghe, 2005)	Cognitive-Behavior Therapy (CBT; Molenaar et al., 2009)
55.	Durham, R. C., Murphy, T., Allan, T., Richard, K., Treliving, L. R., & Fenton, G. W., et al. (1994). Cognitive therapy, analytic psychotherapy and anxiety management training for generalised anxiety disorder. <i>Br J Psychiatry</i> , 165, 315-323. http://www.ncbi.nlm.nih.gov/pubmed/7994500	Generalized anxiety disorder (GAD)	Analytic-based psychotherapy (no specific reference)	Cognitive Therapy / Anxiety Management training (Beck & Emery, 1985)
56.	Emmelkamp, P. M. G., Benner, A., Kuipers, A., Feiertag, G. A., Koster, H. C., & Van Apeldoorn, F. J., et al. (2006). Comparison of brief dynamic and cognitive-behavioural therapies in avoidant personality disorder. <i>Br J Psychiatry</i> , 189, 60-64. http://www.ncbi.nlm.nih.gov/pubmed/16816307	Avoidant personality disorder	Brief Dynamic Therapy (BDT; Malan, 1979; also reference to Luborsky, 1984)	CBT (Beck & Freeman, 1990; Emmelkamp et al, 1992)
57.	Fairburn, G., Kirk, J., Connor, M. O., & Cooper, P. J. (1986). A comparison of two psychological	Bulimia	Short-term Focal Psychotherapy (Rosen,	CBT (Fairburn, 1985)

	treatments for bulimia. <i>Behaviour research and therapy</i> , 24(6), 629–643. http://www.sciencedirect.com/science/article/pii/0005796786900586		1979)	
58.	Faramarzi, M., Azadfallah, P., Book, H. E., Tabatabaei, K. R., Taheri, H., & Shokri-Shirvani, J. (2013). A randomized controlled trial of brief psychoanalytic psychotherapy in patients with functional dyspepsia. <i>Asian journal of psychiatry</i> , 6(3), 228–34. http://www.ncbi.nlm.nih.gov/pubmed/23642981 Faramarzi, M., & Book, H. E. (2015). The Effect of Psychotherapy in Improving Physical and Psychiatric Symptoms in Patients with Functional Dyspepsia, <i>Iran Journal of Psychiatry</i> , 43–49. http://www.ncbi.nlm.nih.gov/pubmed/26005480	Functional dyspepsia	Brief psychodynamic therapy (Book, 1998) based on Luborsky (1984) + standard medical therapy	TAU (medical therapy)
59.	Fonagy, P., Rost, F., Carlyle, J.-A., Mcpherson, S., Thomas, R., Fearon, R. M. P., ... Taylor, D. (2015). Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression : the Tavistock Adult Depression Study (TADS). <i>World Psychiatry</i> , 14(3), 312-321 http://www.ncbi.nlm.nih.gov/pubmed/26407787	Depression (treatment resistant)	Long-Term Psychoanalytic Psychotherapy (LTTP; Taylor, 2015)	TAU
60.	Fonagy, P., Slead, M., & Baradon, T. (2016). Randomized Controlled Trial of Parent-Infant Psychotherapy for Parents With Mental Health Problems and Young Infants. <i>Infant Mental Health Journal</i> , 37(2), 97-114 http://doi.wiley.com/10.1002/imhj.21553	Parents with mental health problems	Parent-infant psychotherapy (PIP)	Standard care
61.	Gallagher-Thompson, D., & Steffen, A. M. (1994). Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. <i>J Consult Clin Psychol</i> , 62, 543-549. http://www.ncbi.nlm.nih.gov/pubmed/8063980	Depression (among family caregivers)	Brief Psychodynamic Therapy (Mann, 1973)	CBT (Beck et al., 1979; Lewinsohn et al, 1985; Gallagher et al 1981)
62.	Gallagher, D. E., & Thompson, L. W. (1982). Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. <i>Psychotherapy: Theory, Research, and Practice</i> , 482-490. http://psycnet.apa.org/journals/pst/19/4/482	Depression	Brief relational/insight psychotherapy (Bellak & Small, 1965)	Behavioral therapy / Cognitive therapy (Lewinsohn et al, 1976; Gallagher et al., 1981)
63.	Garner, D. M., Rockert, W., Davis, R., Garner, M. V., Olmsted, M. P., & Eagle, M., et al. (1993). Comparison of cognitive-behavioral and supportive-expressive therapy for bulimia nervosa. <i>Am J Psychiatry</i> , 150, 37-46. http://www.ncbi.nlm.nih.gov/pubmed/8417578	Bulimia	Supportive-expressive psychotherapy (Luborsky, 1984)	CBT (Fairburn, 1985)
64.	Gelder, M. G., Marks, I. M., & Wolff, H. H. (1967). Desensitization and psychotherapy in the treatment of phobic states: a controlled inquiry. <i>British Journal of Psychiatry</i> , 113(494), 53–81. http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53	Phobic states	Psychotherapy (no reference)	1. Desensitization (Wolpe) 2. Group psychotherapy (Foulkes)

65.	<p>Gersons, B. P., Carlier, I. V, Lamberts, R. D., & van der Kolk, B. a. (2000). Randomized clinical trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. <i>Journal of Traumatic Stress, 13</i>(2), 333–47. http://www.ncbi.nlm.nih.gov/pubmed/10838679</p>	PTSD	Brief eclectic psychotherapy (BEP; Gersons & Carlier, 1994; Manual includes both CBT and PDT components)	Waitlist
66.	<p>Ghorbani, N., Dadsetan, P., Ejei, J., & Motiyan, H. (2000). The consequences of overcoming resistance and emotional disclosure on lymphocyte T-helper and T-suppressor and psychological pathology. <i>Journal of Psychology (Persian), 3</i>, 368–89.</p>	Students	ISTDP (Davanloo, 1990)	Simple verbal disclosure groups (SVD)
67.	<p>Giesen-Bloo, J., van Dyck, R., Spinhoven, P., van Tilburg, W., Dirksen, C., & van Asselt, T., et al. (2006). Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. <i>Arch Gen Psychiatry, 63</i>, 649-658. http://www.ncbi.nlm.nih.gov/pubmed/16754838</p> <p>van Asselt, A. D. I., Dirksen, C. D., Arntz, A., Giesen-Bloo, J. H., van Dyck, R., Spinhoven, P., ... Severens, J. L. (2008). Out-patient psychotherapy for borderline personality disorder: cost-effectiveness of schema-focused therapy v. transference-focused psychotherapy. <i>The British Journal of Psychiatry : The Journal of Mental Science, 192</i>(6), 450–7. http://www.ncbi.nlm.nih.gov/pubmed/18515897</p>	Borderline personality disorder	Transference focused psychotherapy (TFP; Clarkin, Yeomans, & Kernberg, 1999; Yeomans, Clarkin, & Kernberg, 2002)	Schema-focused therapy (Young, 1994; Young et al, 2003)
68.	<p>Gilboa-Schechtman, E., Foa, E. B., Shafran, N., Aderka, I. M., Powers, M. B., Rachamim, L., ... Apter, A. (2010). Prolonged exposure versus dynamic therapy for adolescent PTSD: a pilot randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry, 49</i>(10), 1034–42. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&tool=pmcentrez&rendertype=abstract</p>	PTSD (adolescents)	Time-limited dynamic therapy (TLDP-A; Shefler et al, 1995)	Prolonged exposure (Foa)
69.	<p>Gowers, S., Norton, K., Halek, C., & Crisp, A. H. (1994). Outcome of outpatient psychotherapy in a random allocation treatment study of anorexia nervosa. <i>Int J Eat Disord, 15</i>, 165-177. http://www.ncbi.nlm.nih.gov/pubmed/8173562</p>	Anorexia (adolescents and young women)	Short term individual + family interviews (Koss & Butcher, 1986)	Assessment only
70.	<p>Gregory, R. J., Chlebowski, S., Kang, D., Remen, A. L., Soderberg, M. G., & Stepkovitch, J., et al. (2008). A controlled trial of psychodynamic psychotherapy for co-occurring borderline personality disorder and alcohol use disorder. <i>Psychotherapy: Theory, Research, Practice, Training, 45</i>, 28-41. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28</p>	Borderline personality disorder + alcohol use disorder	Dynamic deconstructive therapy (Gregory & Remen)	TAU
71.	<p>Grothe, C., Bering, R., SpieB, R., Luneburg, E., & Fischer, G. (2003). Mehrdimensionale Psychodynamische Traumaterapie (MPTT) Forschungsergebnisse zur Standardversion. <i>Zeitschrift fur Psychotraumatologie,</i></p>	PTSD	Mehrdimensionale Psychodynamische Traumaterapie (MPTT; Fischer, 2000)	1. EMDR (Shapiro, 1995; 1998) 2. Waitlist control (?)

	<i>Psychotherapiewissenschaft und Psychologische Medizin</i> 1(2), 27–43.			
72.	Gunderson, J. G., Frank, A. F., Katz, H. M., Vannicelli, M. L., Frosch, J. P., & Knapp, P. H., et al. (1984). Effects of psychotherapy in schizophrenia: II. Comparative outcome of two forms of treatment. <i>Schizophr Bull</i> , 10, 564-598. http://www.ncbi.nlm.nih.gov/pubmed/6151246	Schizophrenia	Exploratory, insight oriented psychotherapy (Adapted from Fromm-Reichmann)	Reality-adaptive, supportive psychotherapy
73.	Guthrie, E., Creed, F., Dawson, D., & Tomenson, B. (1991). A controlled trial of psychological treatment for the irritable bowel syndrome. <i>Gastroenterology</i> , 100, 450-457. http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf Guthrie, E., Creed, F., Dawson, D., & Tomenson, B. (1993). A randomised controlled trial of psychotherapy in patients with refractory Irritable Bowel Syndrome. <i>British Journal of Psychiatry</i> , 163, 315–21. http://bjp.rcpsych.org/content/163/3/315	Irritable bowel syndrome	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	Supportive listening
74.	Guthrie, E., Kapur, N., Mackway-Jones, K., Chew-Graham, C., Moorey, J., & Mendel, E., et al. (2001). Randomised controlled trial of brief psychological intervention after deliberate self poisoning. <i>BMJ</i> , 323, 135-138. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&tool=pmcentrez&rendertype=abstract	Self-poisoning	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU
75.	Guthrie, E., Moorey, J., Margison, F., Barker, H., Palmer, S., & McGrath, G., et al. (1999). Cost-effectiveness of brief psychodynamic-interpersonal therapy in high utilizers of psychiatric services. <i>Arch Gen Psychiatry</i> , 56, 519-526. http://www.ncbi.nlm.nih.gov/pubmed/10359466	Mixed sample	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU
76.	Hall, A., & Crisp, A. H. (1987). Brief psychotherapy in the treatment of anorexia nervosa: Outcome at one year. <i>British Journal of Psychiatry</i> , 151, 185-191. http://www.ncbi.nlm.nih.gov/pubmed/3690108	Anorexia	Combined individual and family psychotherapy (no reference)	Dietary advice
77.	Hamilton, J., Guthrie, E., Creed, F., Thompson, D., Tomenson, B., & Bennett, R., et al. (2000). A randomized controlled trial of psychotherapy in patients with chronic functional dyspepsia. <i>Gastroenterology</i> , 119, 661-669. http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949	Chronic Functional Dyspepsia	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	Supportive therapy (Guthrie, unpub.)
78.	Hardy, G. E., Barkham, M., Shapiro, D. a, Stiles, W. B., Rees, a, & Reynolds, S. (1995). Impact of Cluster C personality disorders on outcomes of contrasting brief psychotherapies for depression. <i>Journal of consulting and clinical psychology</i> , 63(6), 997–1004. http://www.ncbi.nlm.nih.gov/pubmed/8543722	Depression + personality disorder cluster C	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	CBT (Firth & Shapiro, 1985)
79.	Hellerstein, D. J., Rosenthal, R. N., Pinsker, H., Samstag, L. W., Muran, J. C., & Winston, A., et	Personality disorders (cluster C)	ISTDP (Davanloo, 1980)	Brief Adaptive Therapy (Winston?)

	<p>al. (1998). A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. <i>J Psychother Pract Res</i>, 7, 261-271. http://psycnet.apa.org/psycinfo/1998-12001-001</p>			
80.	<p>Hersen, M., Himmelhoch, J. M., Thase, M. E., & Bellack, A. S. (1984). Effects of social skill training, amitriptyline, and psychotherapy in unipolar depressed women. <i>Behavior Therapy</i>, 15(1), 21–40. http://linkinghub.elsevier.com/retrieve/pii/S005789484800398</p>	Depression	Time-limited dynamic therapy (no reference) + placebo	<p>1. Social skill training + placebo</p> <p>2. Amitriptyline</p>
81.	<p>Hoffart, A., & Martinsen, E. W. (1990). Exposure-based integrated vs. Pure psychodynamic treatment of agoraphobic inpatients. <i>Psychotherapy: Theory, Research, Practice, Training</i>, Vol 27(2), 210–218. http://psycnet.apa.org/psycinfo/1990-29285-001</p>	Agoraphobia	Integrative behavioural-psychodynamic treatment	Psychodynamic oriented group therapy
82.	<p>Høglend, P., Amlo, S., Marble, A., Bøggwald, K. P., Sørbye, O., & Sjaastad, M. C., et al. (2006). Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental study of transference interpretations. <i>Am J Psychiatry</i>, 163, 1739-1746. http://www.ncbi.nlm.nih.gov/pubmed/17012684</p> <p>Høglend, P., Bøggwald, K.-P., Amlo, S., Marble, A., Ulberg, R., Sjaastad, M. C., ... Johansson, P. (2008). Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? <i>The American Journal of Psychiatry</i>, 165(6), 763–71. http://www.ncbi.nlm.nih.gov/pubmed/18413707</p>	Mixed sample	Psychodynamic psychotherapy with transference interpretation (Høglend, 1995)	Psychodynamic psychotherapy without transference interpretation (Høglend, 1995)
83.	<p>Huber, D., Henrich, G., Gastner, J., & Klug, G. (2012). Must All Have Prizes? The Munich Psychotherapy Study. In <i>Psychodynamic Psychotherapy Research</i>. http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1</p> <p>Huber, D., Huber, D., Zimmermann, J., Henrich, G., & Klug, G. (2012). Comparison of cognitive-behaviour therapy with psychoanalytic and psychodynamic therapy for depressed patients – A three-year follow-up study. <i>Zeitschrift für Psychosomatic</i>, (58), 299–316. http://www.ncbi.nlm.nih.gov/pubmed/22987495</p> <p>Huber, D., Henrich, G., Clarkin, J., & Klug, G. (2013). Psychoanalytic versus psychodynamic therapy for depression: a three-year follow-up study. <i>Psychiatry</i>, 76(2), 132–49. http://www.ncbi.nlm.nih.gov/pubmed/23631544</p>	Depression	Psychoanalysis (non-manualized)	<p>1. LTPP (non-manualized)</p> <p>2. CBT (non-manualized; patients not randomly assigned to this group)</p>

84.	<p>Hubschmid, M., Aybek, S., D, M., Maccaferri, G. E., D, M., Chocron, O., ... D, M. (n.d.). Efficacy of brief interdisciplinary psychotherapeutic intervention for motor conversion disorder and nonepileptic attacks. <i>General Hospital Psychiatry</i>, 37(5), 448–455. doi:10.1016/j.genhosppsy.2015.05.007 http://dx.doi.org/10.1016/j.genhosppsy.2015.05.007</p>	Motor conversion disorder and nonepileptic seizures	Interdisciplinary psychotherapeutic intervention (Psychodynamic-Interpersonal therapy based on Guthrie, 1999)	Standard care
85.	<p>Jantschek, G., Zeitz, M., Wirsching, M., Klor, H. U., Studt, H. H., Rasenack, J., ... Disease, T. G. S. G. on P. I. in C. (1999). Effect of psychotherapy on the course of Crohn's disease. <i>Psychosomatic Intervention in CD</i>, 1289–1296. http://www.ncbi.nlm.nih.gov/pubmed/9930393</p> <p>Keller, W., Pritsch, M., Von Wietersheim, J., Scheib, P., Osborn, W., & Balck, F., et al. (2004). Effect of psychotherapy and relaxation on the psychosocial and somatic course of Crohn's disease: main results of the German Prospective Multicenter Psychotherapy Treatment study on Crohn's Disease. <i>J Psychosom Res</i>, 56, 687-696. http://www.ncbi.nlm.nih.gov/pubmed/15193965</p>	Crohn's disease	PDT+relaxation+drug treatment (no manual was used but reference to Luborsky, 1984)	Drug treatment
86.	<p>Johansson, R., Björklund, M., Hornborg, C., Karlsson, S., Hesser, H., Ljótsson, B., ... Andersson, G. (2013). Affect-focused psychodynamic psychotherapy for depression and anxiety through the Internet: a randomized controlled trial. <i>PeerJ</i>, 1, e102. https://peerj.com/articles/102</p>	Depression and/or anxiety	IPDT (Affect-focused; based on Frederick, 2009)	Online therapist support and clinical monitoring
87.	<p>Johansson, R., Ekbladh, S., Hebert, A., Lindström, M., Möller, S., Petitt, E., ... Andersson, G. (2012). Psychodynamic Guided Self-Help for Adult Depression through the Internet: A Randomised Controlled Trial. (A. Bruce, Ed.) <i>PLoS ONE</i>, 7(5), http://dx.plos.org/10.1371/journal.pone.0038021</p>	Depression	IPDT (SUBGAP; based on Silverberg, 2005)	Psychoeducation + support
88.	<p>Jørgensen, C. R., Freund, C., Bøye, R., Jordet, H., Andersen, D., & Kjølbye, M. (2012). Outcome of mentalization-based and supportive psychotherapy in patients with borderline personality disorder: a randomized trial. <i>Acta psychiatrica Scandinavica</i>, 1–13. http://www.ncbi.nlm.nih.gov/pubmed/22897123</p>	Borderline personality disorder	Mentalization-Based Therapy (MBT; Bateman & Fonagy, 2001)	Supportive group therapy
89.	<p>Klein, D. F., Zitrin, C. M., Woerner, M. G., & Ross, D. C. (1983). Treatment of Phobias. II. Behavior therapy and supportive psychotherapy: are there any specific ingredients? <i>Archives of General Psychiatry</i>, 139–145. http://www.ncbi.nlm.nih.gov/pubmed/6130751</p>	Phobias	Supportive therapy (dynamically oriented; no reference)	Behavior therapy (no reference)
90.	<p>Kleinman, P. H., Woody, G. E., Todd, T. C., Millman, R. B., Kang, S. Y., & Kemp, J., et al. (1990). Crack and cocaine abusers in outpatient psychotherapy. <i>NIDA Res Monogr</i>, 104, 24-35.</p>	Cocaine abuse	Supportive-Expressive Therapy (Luborsky, 1984)	Family therapy + para professionally led group therapy

91.	<p>Knekt, P., & Lindfors, O. (2004). A randomized trial of the effect of four forms of psychotherapy on depressive and anxiety disorders. Helsinki: Kela. https://helda.helsinki.fi/handle/10250/3315</p> <p>Knekt, P., Lindfors, O., Ske, L. S. Ä., & Virtala, E. S. A. (2013). Randomized trial on the effectiveness of long- and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. <i>Nordic journal of psychiatry</i>, 67(1), 59-68. http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910</p> <p>Knekt, P., Virtala, E., Härkänen, T., Vaarama, M., Lehtonen, J., & Lindfors, O. (2016). The outcome of short- and long-term psychotherapy 10 years after start of treatment. <i>Psychological Medicine</i>, 1-14. http://www.journals.cambridge.org/abstract/S0033291715002718</p> <p>Lindfors, O., Knekt, P., Virtala, E., & Laaksonen, M. a. (2012). The effectiveness of solution-focused therapy and short- and long-term psychodynamic psychotherapy on self-concept during a 3-year follow-up. <i>The Journal of Nervous and Mental Disease</i>, 200(11), 946-53. http://www.ncbi.nlm.nih.gov/pubmed/23124178</p> <p>Lindfors, O., Knekt, P., Heinonen, E., Härkänen, T., Virtala, E., & The Helsinki Psychotherapy Study Group. (2015). The effectiveness of short- and long-term psychotherapy on personality functioning during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 173, 31-8. http://www.ncbi.nlm.nih.gov/pubmed/25462393</p> <p>Maljanen, T., Knekt, P., Lindfors, O., Virtala, E., Tillman, P., & Härkänen, T. (2016). The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 190(July), 254-263. http://www.ncbi.nlm.nih.gov/pubmed/22611089</p>	Depression and/or anxiety	STPP (Malan, 1976; Sifneos, 1978)	<p>1. Solution-focused therapy (de Shazer et al, 1986)</p> <p>2. LTPP (Gabbard, 2004)</p>
92.	<p>Knijnik, D. Z., Blanco, C., Salum, G. A., Moraes, C. U., Mombach, C., Almeida, E., ... Eizirik, C. L. (2008). A pilot study of clonazepam versus psychodynamic group therapy plus clonazepam in the treatment of generalized social anxiety disorder. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 23(8), 567-74. http://www.ncbi.nlm.nih.gov/pubmed/18774274</p>	Social anxiety disorder (SAD)	Psychodynamic group therapy (Malan, 1976) + clonazepam	Clonazepam
93.	<p>Knijnik, D. Z., Kapczinski, F., Chachamovich, E., Margis, R., & Eizirik, C. L. (2004). Psychodynamic group treatment for generalized social phobia. <i>Revista Brasileira de Psiquiatria (São Paulo, Brazil : 1999)</i>, 26(2), 77-81. http://www.ncbi.nlm.nih.gov/pubmed/15517057</p>	Social anxiety disorder (SAD)	Psychodynamic group therapy (Malan, 1976)	Credible placebo group

94.	Kompoliti, K., Wilson, B., Stebbins, G., Bernard, B., & Hinson, V. (2013). Immediate vs. delayed treatment of psychogenic movement disorders with short-term psychodynamic psychotherapy: Randomized clinical trial. <i>Parkinsonism & related disorders</i> , 7–10. http://www.ncbi.nlm.nih.gov/pubmed/24120952	Psychogenic movement disorders	ISTDP (Davanloo, 1980)	Neurological observation and support
95.	Kool, S., Dekker, J., & Duijsens, I. (2003). Efficacy of combined therapy and pharmacotherapy for depressed patients with or without personality disorders. <i>Harvard review of psychiatry</i> , 133–142. http://informahealthcare.com/doi/abs/10.1080/10673220303950	Depression	STDP (de Jonghe, 2005) + pharmacotherapy	Pharmacotherapy
96.	Kornblith, S., Rehm, L., W, O. M., & Lamparski, D. M. (1983). The contribution of self-reinforcement training and behavioral assignments to the efficacy of self-control therapy for depression. <i>Cogn Ther Res</i> , 7, 499-528. http://link.springer.com/10.1007/BF01172888	Depression	Psychodynamic group therapy (Wolf, 1965; Bernard & Klein, 1977)	Self-reinforcement training and behavioral assignments (Rehm, 1977)
97.	Kramer, U., de Roten, Y., Perry, J. C., & Despland, J.-N. (2013). Change in defense mechanisms and coping patterns during the course of 2-year-long psychotherapy and psychoanalysis for recurrent depression: a pilot study of a randomized controlled trial. <i>The Journal of nervous and mental disease</i> , 201(7), 614–20. http://www.ncbi.nlm.nih.gov/pubmed/23817160	Depression (recurrent)	Supportive-Expressive Therapy (Luborsky, 1984) All patients were also given pharmacological treatment	1. Cognitive therapy (Beck, 1995) 2. Psychoanalysis (Robertson, 2002). 3. Clinical management (Csank, 2002; Novalis et al., 1993).
98.	Lanza, M. L., Anderson, J., Boisvert, C. M., LeBlanc, A., Fardy, M., & Steel, B., et al. (2002). Assaultive behavior intervention in the Veterans Administration: psychodynamic group psychotherapy compared to cognitive behavior therapy. <i>Perspect Psychiatr Care</i> , 38, 89-97. http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract	Male veterans with history of assault	Psychodynamic group psychotherapy (no specific reference)	Cognitive behavior group therapy (No specific reference)
99.	LaPonite, K., & Rimm, D. C. (1980). Cognitive, assertive, and insight-oriented group therapies in the treatment of reactive depression in women. <i>Psychotherapy: Theory, Research & Practice</i> , 17(3), 312–321. http://psycnet.apa.org/journals/pst/17/3/312	Depression	Insight-oriented group therapy (No reference)	1. Cognitive group therapy (Beck, 19767) 2. Assertiveness training (no reference)
100.	Lau, M., & Kristensen, E. (2007). Outcome of systemic and analytic group psychotherapy for adult women with history of intrafamilial childhood sexual abuse: a randomized controlled study. <i>Acta Psychiatr Scand</i> , 116, 96-104. http://www.ncbi.nlm.nih.gov/pubmed/17650270	Women with childhood sexual abuse	Analytic group therapy (Foulkes, 1986)	Systemic group therapy (Holme, 1999)
101.	Lawson, D. M. (2010). Comparing cognitive behavioral therapy and integrated cognitive behavioral therapy/psychodynamic therapy in group treatment for partner violent men. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 47(1), 122–133. http://doi.apa.org/getdoi.cfm?doi=10.1037/a	Violent men	Integrative CBT-psychodynamic group therapy (Strupp & Binder, 1984)	CBT (Lawson, 2001)

	0018841			
102.	<p>Leichsenring, F., Masuhr, O., Jaeger, U., Rabung, S., Dally, A., Dumpelmann, M., ... Streeck, U. (2016). Psychoanalytic-Interactional Therapy versus Psychodynamic Therapy by Experts for Personality Disorders: A Randomized Controlled Efficacy-Effectiveness Study in Cluster B Personality Disorders. <i>Psychotherapy and Psychosomatics</i>, 71–80. http://www.karger.com/?doi=10.1159/000441731</p>	Personality disorders (cluster B)	Psychoanalytic-Interactional Therapy (Streeck, 2016)	Psychodynamic therapy by experts
103.	<p>Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2013). Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>American Journal of Psychiatry</i>, 170(7), 759-767. http://www.ncbi.nlm.nih.gov/pubmed/23680854</p> <p>Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2014). Long-Term Outcome of Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder. <i>American Journal of Psychiatry</i>, 171(10), 1074-1082. http://www.ncbi.nlm.nih.gov/pubmed/25016974</p>	Social anxiety disorder (SAD)	Supportive-Expressive Therapy (Luborsky, 1984)	CBT (Clark & Wells, 1995)
104.	<p>Leichsenring, F., Salzer, S., Jaeger, U., Kächele, H., Kreische, R., & Leweke, F., et al. (2009). Short-Term Psychodynamic Psychotherapy and Cognitive-Behavioral Therapy in Generalized Anxiety Disorder: A Randomized, Controlled Trial. <i>Am J Psychiatry</i>, 166, 875-881. http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;atab=7</p> <p>Salzer, S., Winkelbach, C., Leweke, F., Leibing, E., & Leichsenring, F. (2011). Generalized Anxiety Disorder: 12-Month Follow-Up. <i>Canadian Journal of Psychiatry</i>, 56(8), 1-10. http://www.ncbi.nlm.nih.gov/pubmed/24229481</p>	Generalized anxiety disorder (GAD)	Supportive-Expressive Therapy (Luborsky, 1984)	CBT (Borcovec & Rucio, 2001, Brown, 1993)
105.	<p>Lemma, A., & Fonagy, P. (2013). Feasibility study of a psychodynamic online group intervention for depression. <i>Psychoanalytic Psychology</i>, 30(3), 367–380. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239</p>	Depression	Online Group Dynamic Interpersonal Therapy (OLDIT; Fonagy & Target, 2008)	Internet discussion group
106.	<p>Lerner, A., Sigal, M., Bacalu, A., & Gelkopf, M. (1992). Short term versus long term psychotherapy in opioid dependence: A pilot study. <i>Israeli Journal of Psychiatry and Related Sciences</i>, 29(2), 114–9. http://www.ncbi.nlm.nih.gov/pubmed/1526756</p>	Opiate dependence	Short-Term Dynamic Psychotherapy (Luborsky, 1989; Mann, 1973; Davanloo, 1978)	Long-Term Dynamic Psychotherapy (Karasu, 1989)
107.	<p>Levy Berg, A., Sandell, R., & Sandahl, C. (2009). Affect-focused body psychotherapy in patients with generalized anxiety disorder: Evaluation of an integrative method. <i>Journal of Psychotherapy Integration</i>, 19(1), 67–85.</p>	Generalized anxiety disorder (GAD)	Affect-Focused Body Psychotherapy (Monsen & Monsen, 2000)	TAU

	http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324			
108.	Lieberman, R. P., & Eckman, T. (1981). Behavior therapy vs insight-oriented therapy for repeated suicide attempters. <i>Archives of General Psychiatry, 38</i> (10), 1126–1130. http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007	Repeated suicide attempts	Insight-Oriented Psychotherapy	Behavior Therapy
109.	Lindauer, R. J. L., Gersons, B. P. R., Van Meijel, E. P. M., Blom, K., Carlier, I. V. E., Vrijlandt, I., & Olff, M. (2005). Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: Randomized clinical trial. <i>Journal of Traumatic Stress, 18</i> (3), 205–212. http://doi.wiley.com/10.1002/jts.20029	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al., 2004; Manual includes both CBT and PDT components)	Waitlist
110.	Linnet, J., & Jemec, G. B. (2001). Anxiety level and severity of skin condition predicts outcome of psychotherapy in atopic dermatitis patients. <i>Int J Dermatol, 40</i> , 632–636. http://www.ncbi.nlm.nih.gov/pubmed/11737422	Atopic dermatitis (anxiety)	Brief Dynamic Psychotherapy (Malan, 1992)	Control (TAU?)
111.	Lipsitz, J. D., Gur, M., Vermes, D., Petkova, E., Cheng, J., Miller, N., ... Fyer, A. J. (2008). A randomized trial of interpersonal therapy versus supportive therapy for social anxiety disorder. <i>Depression and Anxiety, 25</i> (6), 542–553. http://doi.wiley.com/10.1002/da.20364	Social anxiety disorder (SAD)	Psychodynamic Supportive Therapy (Lipsitz et al., 2006)	Interpersonal Therapy (Klerman et al., 1984; Weissman et al., 2000)
112.	Lock, J., Grange, D. Le, & Agras, W. (2010). Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. <i>Archives of General Psychiatry, 67</i> (10), 1025–1032. http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf	Anorexia	Adolescent Focused Therapy (AFT, Robin et al.)	Treatments Family-Based Treatment (FBT; Locke et al., 2001)
113.	Lorentzen, S., Ruud, T., Fjeldstad, A., & Høglend, P. (2013). Comparison of short- and long-term dynamic group psychotherapy: randomised clinical trial. <i>The British journal of psychiatry : the journal of mental science.</i> http://www.ncbi.nlm.nih.gov/pubmed/24029539 Lorentzen, S., Ruud, T., Fjeldstad, A., & Høglend, P. a. (2014). Personality disorder moderates outcome in short- and long-term group analytic psychotherapy: A randomized clinical trial. <i>The British Journal of Clinical Psychology / the British Psychological Society, 1–18.</i> http://www.ncbi.nlm.nih.gov/pubmed/25178520 Fjeldstad, A., Høglend, P., & Lorentzen, S. (2015). Patterns of Change in Interpersonal Problems During and After Short-term and Long-term Psychodynamic Group Therapy: A Randomized Clinical Trial. <i>Psychotherapy Research, 33</i> 07(November), 1–12. http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357	Anxiety and personality disorders	Long-Term Psychodynamic Group Therapy (Lorentzen, 2013)	Short-Term Psychodynamic Group Therapy (Lorentzen, 2013)

114.	<p>Ludwig, G., Krenz, S., Zdrojewski, C., Bot, M., Rousselle, I., Stagno, D., ... Stiefel, F. (2013). Psychodynamic interventions in cancer care I: psychometric results of a randomized controlled trial. <i>Psycho-oncology</i>, 23, 65-74. http://www.ncbi.nlm.nih.gov/pubmed/23983096</p>	Mixed depression and anxiety (in cancer patients)	Brief Dynamic Psychotherapy (Mann, 1973; Davanloo, 1994)	Waitlist
115.	<p>Maina, G., Forner, F., & Bogetto, F. (2005). Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. <i>Psychother Psychosom</i>, 74, 43-50. http://www.ncbi.nlm.nih.gov/pubmed/15627856</p>	Depression (mild to moderate)	Brief Dynamic Psychotherapy (BDP; Malan, 1979)	<p>1. Brief Supportive Psychotherapy (BSP; Novalis et al., 1993))</p> <p>2. Waitlist</p>
116.	<p>Maina, G., Rosso, G., Crespi, C., & Bogetto, F. (2007). Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. <i>Psychotherapy and Psychosomatics</i>, 76(5), 298–305. http://www.ncbi.nlm.nih.gov/pubmed/17700050</p> <p>Maina, G., Rosso, G., & Bogetto, F. (2009). Brief dynamic therapy combined with pharmacotherapy in the treatment of major depressive disorder: long-term results. <i>Journal of Affective Disorders</i>, 114(1-3), 200–7. http://www.ncbi.nlm.nih.gov/pubmed/18728001</p>	Depression	Brief Dynamic Psychotherapy (BDP; Malan, 1963; 1976) + medication	Brief Supportive Psychotherapy (BSP; Novalis et al., 1993) + medication
117.	<p>Maina, G., Rosso, G., Rigardetto, S., Chiadò Piat, S., & Bogetto, F. (2010). No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. <i>Psychotherapy and Psychosomatics</i>, 79(5), 295–302. http://www.ncbi.nlm.nih.gov/pubmed/20616624</p>	OCD + depression	Brief Dynamic Psychotherapy (BDP; Malan, 1979) + medication	Medication only
118.	<p>Marmar, C. R., Horowitz, M. J., Weiss, D. S., Wilner, N. R., & Kaltreider, N. B. (1988). A controlled trial of brief psychotherapy and mutual-help group treatment of conjugal bereavement. <i>Am J Psychiatry</i>, 145, 203-209. http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf</p>	Bereavement	Brief Dynamic Psychotherapy (BDP; Malan, Sifneos; Mann)	Mutual Self-Help Groups
119.	<p>Martini, B., Rosso, G., Chiodelli, D. F., De Cori, D., & Maina, G. (2011). Brief dynamic therapy combined with pharmacotherapy in the treatment of panic disorder with concurrent depressive symptoms. <i>Clinical Neuropsychiatry</i>, 8(3), 204–21. http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf</p>	Panic disorder + depression	Brief Dynamic Psychotherapy (BDP; Malan, 1963) + medication	Brief Supportive Psychotherapy (BSP) + medication
120.	<p>McCallum, M., & Piper, W. E. (1990). A controlled study of effectiveness and patient suitability for short-term group psychotherapy. <i>Int J Group Psychother</i>, 40, 431-452. http://www.ncbi.nlm.nih.gov/pubmed/2228349</p>	Mixed sample	Short-term psychoanalytically oriented group psychotherapy (STG; manual mentioned but no specific reference)	Waitlist
121.	<p>McLean, P., & Hakstian, A. (1979). Clinical depression: comparative efficacy of outpatient</p>	Depression	Short-Term Psychotherapy (Marmor,	1. Relaxation therapy (no reference)

	treatments. <i>J Consult Clin Psychol</i> , 47, 818-836. http://www.ncbi.nlm.nih.gov/pubmed/389965		1975; Wolberg, 1967)	2. Behavior therapy (McLean, 1976) 3. Drug therapy
122.	McMain, S. F., Links, P. S., Gnam, W. H., Guimond, T., Cardish, R. J., & Korman, L., et al. (2009). A randomized trial of dialectical behavior therapy versus general psychiatric management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1365-1374. http://www.ncbi.nlm.nih.gov/pubmed/19755574 McMain, S., Guimond, T., & Streiner, D. (2012). Dialectical Behavior Therapy Compared With General Psychiatric Management for Borderline Personality Disorder: Clinical Outcomes and Functioning Over a 2-Year Follow-Up. <i>American Journal of Psychiatry</i> , 650-661. http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&RelatedWidgetArticles=true	Borderline personality disorder	General psychiatric management, dynamically informed therapy (Gunderson, 1984)	Dialectical Behavior Therapy (DBT; Linehan et al., 1991)
123.	Meyer, A. E., & Bolz, W. (1981). The Hamburg short psychotherapy comparison experiment. Foreword and introduction. <i>Psychother Psychosom</i> , 35, 81-84. http://www.karger.com/doi/10.1159/000287484	Mixed sample	Psychodynamic conflict-centered time-limited psychotherapy (no reference)	Waitlist
124.	Miller, L. C., Barrett, C. L., Hampe, E., & Noble, H. (1972). Comparison of reciprocal inhibition, psychotherapy, and waiting list control for phobic children. <i>Journal of Abnormal Psychology</i> , 79(3), 269-279. http://psycnet.apa.org/journals/abn/79/3/269	Phobia (children)	Play psychotherapy (Lippman, 1956)	Reciprocal inhibition
125.	Milrod, B., Chambless, D. L., Gallop, R., Busch, F. N., Schwalberg, M., McCarthy, K. S., ... Barber, J. P. (2015). Psychotherapies for Panic Disorder: A Tale of Two Sites. <i>The Journal of Clinical Psychiatry</i> , 10065, 1-9. http://www.psychiatrist.com/jcp/article/pages/2015/aheadofprint/14m09507.aspx	Panic disorder	Panic-Focused Psychodynamic Psychotherapy (PFPP; Milrod et al, 2001)	1. CBT (Craske, 2000) 2. Applied relaxation (Öst & Westling, 1995)
126.	Milrod, B., Leon, A. C., Busch, F., Rudden, M., Schwalberg, M., & Clarkin, J., et al. (2007). A randomized controlled clinical trial of psychoanalytic psychotherapy for panic disorder. <i>Am J Psychiatry</i> , 164, 265-272. http://www.ncbi.nlm.nih.gov/pubmed/17267789	Panic disorder	Panic-Focused Psychodynamic Psychotherapy (PFPP; Milrod et al, 1997)	Applied relaxation (Öst & Westling, 1995)
127.	Monsen, K., & Monsen, J. T. (2000). Chronic Pain and Psychodynamic Body Therapy: A Controlled Outcome Study. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 37(3), 257-269. http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2000-02713-007	Chronic pain	Psychodynamic body therapy (Monsen & Monsen, 1999)	TAU
128.	Montag, C., Haase, L., Seidel, D., Bayerl, M., Gallinat, J., Herrmann, U., & Dannecker, K. (2014). A Pilot RCT of Psychodynamic Group Art Therapy for Patients in Acute Psychotic Episodes: Feasibility, Impact on Symptoms and Mentalising Capacity. <i>PLoS One</i> , 9(11),	Schizophrenia	Psychodynamic art therapy (Dannecker, 2010)	TAU

	e112348. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4231093&tool=pmcentrez&rendertype=abstract			
129.	Munroe-Blum, H., & Marziali, E. (1995). A controlled trial of short-term group treatment for borderline personality disorder. <i>J Personal Disord, 9</i> , 190-198. http://psycnet.apa.org/psycinfo/1996-17928-001	Borderline personality disorder	Time-limited group therapy (Marziali & Munroe-Blum, 1995)	Individual psychodynamic psychotherapy (Kernberg, 1975)
130.	Muran, J. C., Safran, J. D., Samstag, L. W., & Winston, A. (2005). Evaluating an alliance-focused treatment for personality disorders. <i>Psychotherapy: Theory, Research, Practice, Training, 42</i> (4), 532–545. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.532 Safran, J. D., Muran, J. C., Samstag, L. W., & Winston, A. (2005). Evaluating alliance-focused intervention for potential treatment failures: A feasibility study and descriptive analysis. <i>Psychotherapy: Theory, Research, Practice, Training, 42</i> (4), 512–531. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.4.512	Personality disorders (mixed; mainly cluster C)	Brief Relational Therapy (BRT; Safran & Muran, 2000)	1. Short-Term Dynamic Psychotherapy (STDP: Pollock, Flegenheimer, & Kaufman, 1992) 2. Cognitive– behavioral therapy (CBT; Turner & Muran, 1992; Beck, Freeman, & Associates, 1990)
131.	Nader-Mohammadi Moghadam, M. Atef-Vahid, M. K., Asgharnejad-Farid, A. A., Shabani, A., & Lavasni, F. (2015). Effectiveness of Short-term Dynamic Psychotherapy versus Sertraline in Treatment of Social Phobia. <i>Iran Journal of Psychiatry, 9</i> (2), 2–6. http://www.ncbi.nlm.nih.gov/pubmed/26288643	Social anxiety disorder (SAD)	Affect Phobia Therapy (APT; McCullough et al., 2003)	1. Sertraline 2. Waitlist
132.	Nijdam, M. J., Gersons, B. P. R., Reitsma, J. B., de Jongh, A., & Olf, M. (2012). Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial. <i>The British Journal of Psychiatry : The Journal of Mental Science, 200</i> (3), 224–31. http://www.ncbi.nlm.nih.gov/pubmed/22322458	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	Eye movement and reprocessing (EMDR; De Jongh et al, 2004)
133.	Ojehagen, A., Berglund, M., Appel, C. P., Andersson, K., Nilsson, B., & Skjaeris, A., et al. (1992). A randomized study of long-term outpatient treatment in alcoholics. Psychiatric treatment versus multimodal behavioural therapy, during 1 versus 2 years of treatment. <i>Alcohol Alcohol, 27</i> , 649-658. http://www.ncbi.nlm.nih.gov/pubmed/1292438	Alcohol dependence	"Psychiatric treatment" based on psychodynamic principles (Luborsky, 1984)	Multimodal behavior therapy (Lazarus, 1981)
134.	Pierloot, R., & Vinck, J. (1978). Differential outcome of short-term dynamic psychotherapy and systematic desensitizations in the treatment of anxious outpatients: A preliminary report. <i>Psychologica Belgica, 18</i> , 87-98. http://psycnet.apa.org/psycinfo/1980-32873-001	Anxiety	Short-Term Dynamic Psychotherapy (Malan, 1963)	Systematic desensitization (Wolpe, 1969)
135.	Piper, W. E., Azim, H. F., McCallum, M., & Joyce, A. S. (1990). Patient suitability and outcome in short-term individual	Mixed sample	STDP (Malan, 1976; Strupp & Binder, 1984)	Waitlist

	psychotherapy. <i>J Consult Clin Psychol</i> , 58, 475-481. http://www.ncbi.nlm.nih.gov/pubmed/2212185			
136.	Piper, W. E., Debbane, E. G., Bienvenu, J. P., & Garant, J. (1984). A comparative study of four forms of psychotherapy. <i>J Consult Clin Psychol</i> , 52, 268-279. http://www.ncbi.nlm.nih.gov/pubmed/6425378	Mixed sample	Short-term individual (STI; Malan, 1976)	1. Short-term group (STG) 2. Long-term individual (LTI) 3. Long-term group (LTG) Non manuals used
137.	Piper, W. E., Joyce, a S., McCallum, M., & Azim, H. F. (1998). Interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i> , 66(3), 558–67. http://www.ncbi.nlm.nih.gov/pubmed/10224738 Piper, W. E., McCallum, M., Joyce, a S., Azim, H. F., & Ogrodniczuk, J. S. (1999). Follow-up findings for interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i> , 67(2), 267–73. http://www.ncbi.nlm.nih.gov/pubmed/10224738	Mixed sample	Interpretive psychotherapy (Piper, Azim, Joyce, & McCallum, 1993)	Supportive psychotherapy (Piper, Azim, Joyce, & McCallum, 1993)
138.	Piper, W. E., McCallum, M., Joyce, A. S., Rosie, J. S., & Ogrodniczuk, J. S. (2001). Patient personality and time-limited group psychotherapy for complicated grief. <i>Int J Group Psychother</i> , 51, 525-552. http://www.ncbi.nlm.nih.gov/pubmed/11582899 Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., Weideman, R., & Rosie, J. S. (2007). Group composition and group therapy for complicated grief. <i>Journal of Consulting and Clinical Psychology</i> , 75(1), 116–25. http://www.ncbi.nlm.nih.gov/pubmed/17295570 Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., & Weideman, R. (2009). Follow-up outcome in short-term group therapy for complicated grief. <i>Group Dynamics: Theory, Research, and Practice</i> , 13(1), 46–58. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830	Complicated grief	Interpretive group therapy (no specific reference)	Supportive group therapy (no specific reference)
139.	Poulsen, A. (1991). Psychodynamic, time-limited group therapy in rheumatic disease—a controlled study with special reference to alexithymia. <i>Psychother Psychosom</i> , 56, 12-23. http://www.ncbi.nlm.nih.gov/pubmed/1891575	Rheumatic disease	Time-limited group therapy (Foulkes, 1975)	Waitlist
140.	Poulsen, S., Lunn, S., Daniel, S. I. F., Folke, S., Mathiesen, B. B., Katznelson, H., & Fairburn, C. G. (2013). A Randomized Controlled Trial of Psychoanalytic Psychotherapy or Cognitive-Behavioral Therapy for Bulimia Nervosa. <i>The American journal of psychiatry</i> , 1–8. http://www.ncbi.nlm.nih.gov/pubmed/24275909	Bulimia	Psychoanalytic Psychotherapy for bulimia nervosa (Lunn & Poulsen, 2012)	CBT (Fairburn, 2008)
141.	Reneses, B., Galián, M., Serrano, R., Figuera,	Borderline personality	Representation focused	TAU

	D., Fernandez Del Moral, A., López-Ibor, J. J., ... Trujillo, M. (2013). A new Time Limited Psychotherapy for BPD: Preliminary Results of a Randomized and Controlled Trial. <i>Actas españolas de psiquiatría</i> , 41(3), 139–48. http://www.ncbi.nlm.nih.gov/pubmed/23803797	disorder	Psychotherapy (PRFP; (Trujillo, 2001; 2010)	
142.	Ridsdale , L., Godfrey, E., Chalder, T., Seed, P., King, M., Wallace, P., & Wessely, S. (2001). Chronic fatigue in general practice: is counselling as good as cognitive behaviour therapy? A UK randomised trial. <i>The British Journal of General Practice : The Journal of the Royal College of General Practitioners</i> , 51(462), 19–24. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&tool=pmcentrez&rendertype=abstract	Chronic fatigue	Counselling “...all counsellors used a psychodynamic approach and all cognitive behaviour therapists used CBT” (p. 20)	CBT
143.	Robin , A L., Siegel, P. T., & Moye, a. (1995). Family versus individual therapy for anorexia: impact on family conflict. <i>The International Journal of Eating Disorders</i> , 17(4), 313–22. http://www.ncbi.nlm.nih.gov/pubmed/7620470 Robin, A. L., Siegel, P. T., Moye, A. W., Gilroy, M., Dennis, A. B., & Sikand, A., et al. (1999). A controlled comparison of family versus individual therapy for adolescents with anorexia nervosa. <i>J Am Acad Child Adolesc Psychiatry</i> , 38, 1482-1489. http://www.ncbi.nlm.nih.gov/pubmed/10596247	Anorexia	Ego-oriented individual therapy (EOIT) (Robin, Siegel, Moye, Koepke, & Tice, 1994)	Behavioral family systems therapy (BFST; Robin et al., 1994, 1996)
144.	Robinson , P., Hellier, J., Barrett, B., Barzdaitiene, D., Bateman, A., Bogaardt, A., ... Fonagy, P. (2016). The NOURISHED randomised controlled trial comparing mentalisation-based treatment for eating disorders (MBT-ED) with specialist supportive clinical management (SSCM-ED) for patients with eating disorders and symptoms of borderline personality disorder. <i>Trials</i> , 17(1), 549. http://doi.org/10.1186/s13063-016-1606-8	Borderline personality disorder + eating disorders	MBT-ED (Bateman & Fonagy, 2009)	Specialist supportive clinical management (SSCM-ED; McIntosh et al., 2006)
145.	Rosenbaum , B., Harder, S., Knudsen, P., Køster, A., Lindhardt, A., Lajer, M., ... Winther, G. (2012). Supportive Psychodynamic Psychotherapy versus Treatment as Usual for First-Episode Psychosis: Two-Year Outcome. <i>Psychiatry</i> , 75(4), 331–41. http://www.ncbi.nlm.nih.gov/pubmed/23244011 Harder, S., Koester, A., Valbak, K., & Rosenbaum, B. (2014). Five-year follow-up of supportive psychodynamic psychotherapy in first-episode psychosis: long-term outcome in social functioning. <i>Psychiatry</i> , 77(2), 155–68. http://www.ncbi.nlm.nih.gov/pubmed/24865198	Schizophrenia	Supportive Psychodynamic Psychotherapy (SPP; Rosenbaum, 2009)	TAU
146.	Rosser , R., Denford, J., Heslop, A., Kinston, W., Macklin, D., & Minty, K., et al. (1983). Breathlessness and psychiatric morbidity in chronic bronchitis and emphysema: a study of psychotherapeutic management. <i>Psychol Med</i> , 13, 93-110.	Breathlessness and psychiatric morbidity (in patients with chronic bronchitis and emphysema)	Analytical psychotherapy (No specific reference)	1. No treatment 2. Nurse support

	http://www.ncbi.nlm.nih.gov/pubmed/6844471			
147.	Rosso, G., Martini, B., & Maina, G. (2012). Brief dynamic therapy and depression severity: A single-blind, randomized study. <i>Journal of affective disorders</i> , 1–6. http://www.ncbi.nlm.nih.gov/pubmed/23174499	Depression	Brief Dynamic Psychotherapy (BDP; Malan, 1976)	Brief Supportive Psychotherapy (BSP; Novalis et al., 1993)
148.	Rossouw, T. I., & Fonagy, P. (2012). Mentalization-based treatment for self-harm in adolescents: a randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 51(12), 1304–1313. http://www.ncbi.nlm.nih.gov/pubmed/23200287	Self-harm (adolescents)	Mentalization-based treatment for adolescents (MBT-A; Rossouw, unpublished)	TAU
149.	Salminen, J. K., Karlsson, H., Hietala, J., Kajander, J., Aalto, S., & Markkula, J., et al. (2008). Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. <i>Psychother Psychosom</i> , 77, 351-357. http://www.ncbi.nlm.nih.gov/pubmed/18701831	Depression	Short-term psychodynamic psychotherapy (STDP; Malan, 1976; Mann, 1973)	Fluoxetine
150.	Salomonsson, B., & Sandell, R. (2011). A randomized controlled trial of mother–infant psychoanalytic treatment: I. Outcomes on self-report questionnaires and external ratings. <i>Infant Mental Health Journal</i> , 32(2), 207–231. http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full	Mothers who worry about child	Mother–infant psychoanalytic treatment (MIP; Norman, 2001; 2004)	Child Health Center Care (TAU)
151.	Salzer, S., Cropp, C., Jaeger, U., Masuhr, O., & Streeck-Fischer, a. (2013). Psychodynamic therapy for adolescents suffering from co-morbid disorders of conduct and emotions in an in-patient setting: a randomized controlled trial. <i>Psychological medicine</i> , 1–10. http://www.ncbi.nlm.nih.gov/pubmed/24229481 Salzer, S., Cropp, C., & Streeck-Fischer, A. (2014). Early Intervention for Borderline Personality Disorder: Psychodynamic Therapy in Adolescents. <i>Zeitschrift Für Psychosomatische Medizin Und Psychotherapie</i> , 60, 368–382. http://doi.org/10.13109/zptm.2014.60.4.368	Mixed sample of adolescents Adolescents with Borderline personality disorder (subsample in secondary study)	Psychoanalytic-interactive method (PiM; Streeck & Leichsenring, 2009; Leichsenring et al., 2010)	Waitlist/TAU
152.	Sanchez, V. C., Lewinsohn, P. M., & Larson, D. W. (1980). Assertion training: effectiveness in the treatment of depression. <i>J Clin Psychol</i> , 36, 526-529. http://www.ncbi.nlm.nih.gov/pubmed/7372826	Depression	“Traditional” Group Therapy (No specific reference)	Assertion training
153.	Sandahl, C., Herlitz, K., Ahlin, G., & Rönnerberg, S. (1998). Time-limited group psychotherapy for moderately alcohol dependent patients: A randomized controlled clinical trial. <i>Psychotherapy Research</i> , 8, 361-378. http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467	Alcohol dependence	Time-limited group psychotherapy (Foulkes, 1964)	Cognitive-behavioural group therapy (Annis, 1986; Marlatt & Gordon, 1985; Sandahl and Rönnerberg, 1990)
154.	Sandahl, C., Lundberg, U., Lindgren, A., Rylander, G., Herlofson, J., Nygren, A., & Asberg, M. (2011). Two forms of group therapy and individual treatment of work-	Depression (work-related)	Focused psychodynamic group therapy (FGT; Sandahl & Lindgren, 2003)	1. Cognitive Group Therapy (no specific reference)

	related depression: a one-year follow-up study. <i>International journal of group psychotherapy</i> , 61(4), 539–55. http://www.ncbi.nlm.nih.gov/pubmed/21985258			2. Comparison condition
155.	Sattel, H., Lahmann, C., Gündel, H., Guthrie, E., Kruse, J., Noll-Hussong, M., ... Henningsen, P. (2011). Brief psychodynamic interpersonal psychotherapy for patients with multisomatoform disorder: randomised controlled trial. <i>The British journal of psychiatry : the journal of mental science</i> , 60–67. http://www.ncbi.nlm.nih.gov/pubmed/22075651	Multisomatoform disorder	Psychodynamic-interpersonal psychotherapy (Guthrie, 2011; Henningsen, 2011)	Enhanced medical care
156.	Saunders, D. G. (1996). Feminist-cognitive-behavioral and process-psychodynamic treatments for men who batter: interaction of abuser traits and treatment models. <i>Violence Vict</i> , 11, 393-414. http://www.ncbi.nlm.nih.gov/pubmed/9210279	Violent men	Process-psychodynamic group therapy (PPT; Browne, Saunders, & Staecker, in press)	Feminist-group CBT (FCBT; Browne, Saunders, & Staecker, in press)
157.	Scheidt, C. E., Waller, E., Endorf, K., Schmidt, S., König, R., Zeeck, A., ... Lacour, M. (2012). Is brief psychodynamic psychotherapy in primary fibromyalgia syndrome with concurrent depression an effective treatment? A randomized controlled trial. <i>General hospital psychiatry</i> . http://www.ncbi.nlm.nih.gov/pubmed/23218844	Fibromyalgia + depression	Psychodynamic-interpersonal psychotherapy (Hobson, 1985; Goldberg et al., 1984; Guthrie, 2013)	TAU
158.	Schnyder, U., & Müller, J. (2011). Brief Eclectic Psychotherapy for PTSD: A Randomized Controlled Trial. <i>The Journal of Clinical ...</i> , 564–566. http://europepmc.org/abstract/MED/21527127	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	Minimal attention control
159.	Shaffer, H. J., LaSalvia, T. A., & Stein, J. P. (1997). Comparing Hatha yoga with dynamic group psychotherapy for enhancing methadone maintenance treatment: a randomized clinical trial. <i>Altern Ther Health Med</i> , 3, 57-66. http://www.ncbi.nlm.nih.gov/pubmed/9210777	Opiate addiction	Dynamic group psychotherapy	Hatha yoga therapy
160.	Shapiro, D. A., Barkham, M., Rees, A., Hardy, G. E., Reynolds, S., & Startup, M., et al. (1994). Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic-interpersonal psychotherapy. <i>J Consult Clin Psychol</i> , 62, 522-534. http://www.ncbi.nlm.nih.gov/pubmed/8063978	Depression	Psychodynamic-interpersonal psychotherapy (Hobson, 1985; Goldberg et al., 1984)	Cognitive Therapy (CT; Beck, Rush, Shaw, and Emery, 1979)
161.	Shapiro, D., & Firth, J. (1987). Prescriptive v. Exploratory Psychotherapy Outcomes of the Sheffield Psychotherapy Project. <i>The British Journal of Psychiatry</i> , 151, 790-799 http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short Shapiro, D. a, & Firth-Cozens, J. (1990). Two-year follow-up of the Sheffield Psychotherapy Project. <i>The British journal of psychiatry</i> , 157,	Deression and anxiety	Exploratory therapy (no specific reference)	Prescriptive therapy (no specific reference)

	389–91. http://www.ncbi.nlm.nih.gov/pubmed/2245269			
162.	Shaw, C. M., Margison, F. R., Guthrie, E. A., & Tomenson, B. (2001). Psychodynamic interpersonal therapy by inexperienced therapists in a naturalistic setting: A pilot study. <i>European Journal of Psychotherapy, Counselling, and Health, 4</i> , 87–101. http://www.tandfonline.com/doi/abs/10.1080/13642530110040091	Mixed sample	Psychodynamic-interpersonal psychotherapy (Hobson, 1985)	Waitlist
163.	Shefler, G., Dasberg, H., & Ben-Shakhar, G. (1995). A randomized controlled outcome and follow-up study of Mann's time-limited psychotherapy. <i>J Consult Clin Psychol, 63</i> , 585-593. http://psycnet.apa.org/journals/ccp/63/4/585	Mixed sample	Time-Limited Psychotherapy (TLP; Mann, 1973)	Waitlist
164.	Simpson, S., Corney, R., Fitzgerald, P., & Beecham, J. (2003). A randomized controlled trial to evaluate the effectiveness and costeffectiveness of psychodynamic counselling for general practice patients with chronic depression. <i>Psychol Med, 33</i> , 229-239. http://www.ncbi.nlm.nih.gov/pubmed/12622302	Depression (chronic)	Psychodynamic counselling (Burton, 1998)	General practice
165.	Sjodin, I., Svedlund, J., Ottosson, J. O., & Dotevall, G. (1986). Controlled study of psychotherapy in chronic peptic ulcer disease. <i>Psychosomatics, 27</i> , 187-91, 195. http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract	Peptic ulcer disease	Short-Term Dynamic Therapy (Malan, 1976)	Medication
166.	Sleed, M., Baradon, T., & Fonagy, P. (2013). New Beginnings for mothers and babies in prison: A cluster randomized controlled trial. <i>Attachment & Human Development, 15</i> (4), 349–67. http://www.ncbi.nlm.nih.gov/pubmed/23550526	High risk mothers (and their babies) in prison	Attachment-based intervention (Baradon, 2006)	Standard health and social care
167.	Sloane, R. B., Staples, F. R., Cristol, A. H., Yorkston, N. J., & Whipple, K. (1975). Short-term analytically oriented psychotherapy versus behavior therapy. <i>American Journal of Psychiatry, 132</i> , 373-377. http://www.ncbi.nlm.nih.gov/pubmed/1119588	Anxiety or personality disorder	Short-Term Analytically Oriented Psychotherapy (no specific reference)	1. Waitlist 2. Behavior Therapy (no specific reference)
168.	Smyrniotis, K. X., & Kirkby, R. J. (1993). Long-term comparison of brief versus unlimited psychodynamic treatments with children and their parents. <i>J Consult Clin Psychol, 61</i> , 1020-1027. http://www.ncbi.nlm.nih.gov/pubmed/8113479	Children with mixed problems	Time-limited brief psychotherapy (Smyrniotis, Kirkby, Smyrniotis, & Picone, 1987)	Time-unlimited psychotherapy (Klein, 1975; Segal, 1978)
169.	Snyder, D. K., Wills, R. M., & Grady-Fletcher, a. (1991). Long-term effectiveness of behavioral versus insight-oriented marital therapy: a 4-year follow-up study. <i>Journal of Consulting and Clinical Psychology, 59</i> (1), 138–41. http://www.ncbi.nlm.nih.gov/pubmed/2002129	Couples with relational problems	Insight-Oriented Marital Therapy (IOMT; no specific reference)	Behavioral Marital Therapy (BMT; no specific reference)
170.	Sørensen, P., Birket-Smith, M., Wattar, U., Buemann, I., & Salkovskis, P. (2010). A	Hypochondriasis	Short-Term Psychodynamic	1. Cognitive-behavioural Therapy (CBT; Salkovskis,

	randomized clinical trial of cognitive behavioural therapy versus short-term psychodynamic psychotherapy versus no intervention for patients with hypochondriasis. <i>Psychol Med</i> , 1-11. http://www.ncbi.nlm.nih.gov/pubmed/20380779		Psychotherapy (No specific reference)	1989; Warwick, 1989; Salkovskis et al. 2003) 2. Waitlist
171.	Stanton, A. H., Gunderson, J. G., Knapp, P. H., Frank, A. F., Vancelli, M. L., & Schnitzer, R., et al. (1984). Effects of psychotherapy in schizophrenia: I. Design and implementation of a controlled study. <i>Schizophr Bull</i> , 10, 520-563. http://www.ncbi.nlm.nih.gov/pubmed/6151245	Schizophrenia (non-chronic)	Expressive, insight-oriented psychotherapy (EIO; Stanton, in press)	Reality-adaptive, supportive therapy (RAS; May, 1968)
172.	Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattheß, H., ... Kruse, J. (2016). Resource activation for treating post-traumatic stress disorder, comorbid symptoms and impaired functioning: a randomized controlled trial in Cambodia. <i>Psychological Medicine</i> , 1–12. http://doi.org/10.1017/S0033291716002592 Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattes, H., ... Kruse, J. (2016). Treating post-traumatic stress disorder by resource activation in Cambodia. <i>World Psychiatry</i> , 15(2), 182–183. http://doi.org/10.1002/wps.20312	PTSD	Resource-Oriented Trauma Therapy (ROTATE; Wöller & Mattheß, 2016)	Waitlist
173.	Svartberg, M., Seltzer, M. H., & Stiles, T. C. (1998). The effects of common and specific factors in short-term anxiety-provoking psychotherapy: a pilot process-outcome study. <i>J Nerv Ment Dis</i> , 186, 691-696. http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx	Mixed sample	Short-term anxiety-provoking psychotherapy (STAPP; Sifneos, 1992)	Non-directive Psychotherapy (NDP; utvecklades specifict för studien)
174.	Svartberg, M., Stiles, T. C., & Seltzer, M. H. (2004). Randomized, controlled trial of the effectiveness of short-term dynamic psychotherapy and cognitive therapy for cluster C personality disorders. <i>Am J Psychiatry</i> , 161, 810-817. http://www.ncbi.nlm.nih.gov/pubmed/15121645	Personality disorder (Cluster C)	Short-Term Dynamic Psychotherapy (STDTP; McCullough Vaillant, 1997)	Cognitive Therapy (CT; Beck & Freeman, 1990)
175.	Svedlund, J., Sjodin, I., Ottosson, J. O., & Dotevall, G. (1983). Controlled study of psychotherapy in irritable bowel syndrome. <i>Lancet</i> , 2, 589-592. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&tool=pmcentrez&rendertype=abstract	IBS	Brief Dynamic Psychotherapy (Malan, 1979)	Treatment as usual (TAU) (medical)
176.	Szapocznik, J., Rio, A., Murray, E., Cohen, R., Scopetta, M., & Rivas-Vazquez, A., et al. (1989). Structural family versus psychodynamic child therapy for problematic Hispanic boys. <i>J.Consult.Clin.Psychol.</i> , 57, 571-578. http://www.ncbi.nlm.nih.gov/pubmed/2794177	Behavioral problems among boys	Individual psychodynamic child therapy (IPCT) (Adams, 1974; Cooper & Wanerman, 1977)	1. Structural Family Therapy (SFT; Minuchin & Fishman, 1981; Minuchin, Montalvo, Guerney, Rosman, & Schuman, 1967) 2. Recreational control condition
177.	Tasca, G., Ritchie, K., Conrad, G., Balfour, L., Gayton, J., & Lybanon, V., et al. (2006). Attachment scales predict outcome in a	Binge eating disorder	Group psychodynamic interpersonal psychotherapy (Tasca et	Group cognitive-behavior therapy (GCBT; Wilfley et al., 1996)

	<p>randomized controlled trial of two group therapies for binge eating disorder: An aptitude by treatment interaction. <i>Psychotherapy Research</i>, 16, 106-121. http://www.tandfonline.com/doi/abs/10.1080/10503300500090928</p>		<p>al., 2002; Strupp & Binder, 1984; Malan 1979)</p>	
178.	<p>Thompson, L. W., Gallagher, D., & Breckenridge, J. S. (1987). Comparative effectiveness of psychotherapies for depressed elders. <i>J Consult Clin Psychol</i>, 55, 385-390. http://www.ncbi.nlm.nih.gov/pubmed/3597953</p>	<p>Depression (among elderly)</p>	<p>Brief psychodynamic therapy (Horowitz & Kaltreider, 1979)</p>	<p>1. Cognitive Therapy (CT; Beck, Rush, Shaw, & Emery, 1979)</p> <p>2. Behavioral Therapy (BT; Gallagher & Thompson, 1981; Lewinsohn, 1974)</p> <p>3. Delayed-treatment control</p>
179.	<p>Thyme, K. E., Sundin, E. C., Stahlberg, G., Lindstrom, B., Eklof, H., & Wiberg, B., et al. (2007). The outcome of short-term psychodynamic art therapy compared to short-term psychodynamic verbal therapy for depressed women. <i>Psychoanalytic Psychotherapy</i>, 21, 250-264. http://www.tandfonline.com/doi/abs/10.1080/02668730701535610</p>	<p>Depression (among women)</p>	<p>Short-Term Psychodynamic Verbal Therapy (Mann, 1973)</p>	<p>Art psychotherapy (no reference)</p>
180.	<p>Trowell, J., Joffe, I., Campbell, J., Clemente, C., Almqvist, F., & Soininen, M., et al. (2007). Childhood depression: a place for psychotherapy. An outcome study comparing individual psychodynamic psychotherapy and family therapy. <i>Eur Child Adolesc Psychiatry</i>, 16, 157-167. http://www.ncbi.nlm.nih.gov/pubmed/17200793</p> <p>Garoff, F. F., Heinonen, K., Pesonen, A.-K., & Almqvist, F. (2012). Depressed youth: treatment outcome and changes in family functioning in individual and family therapy. <i>Journal of Family Therapy</i>, 34(1), 4–23. http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x</p>	<p>Depression (children)</p>	<p>Focused Individual Psychotherapy (Malan & Osmimo, 1992; Davanloo, 1978)</p>	<p>Systems Integrative Family Therapy (Byung-Hall, 1995; Will, 1985)</p>
181.	<p>Trowell, J., Kolvin, I., Weeramanthri, T., Sadowski, H., Berelowitz, M., & Glaser, D., et al. (2002). Psychotherapy for sexually abused girls: psychopathological outcome findings and patterns of change. <i>Br J Psychiatry</i>, 180, 234-247. http://www.ncbi.nlm.nih.gov/pubmed/11872516</p>	<p>Trauma (sexually abused girls)</p>	<p>Individual therapy (Unclear with no reference)</p>	<p>Psychopedagogical /analytic group therapy</p>
182.	<p>Vinnars, B., Barber, J. P., Norén, K., Gallop, R., & Weinryb, R. M. (2005). Manualized supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders: bridging efficacy and effectiveness. <i>Am J Psychiatry</i>, 162, 1933-1940. http://www.ncbi.nlm.nih.gov/pubmed/16199841</p> <p>Vinnars, B., Gallop, R., & Barber, J. P. (2009). Do Personality Problems Improve During Psychodynamic Supportive-Expressive Psychotherapy? Secondary Outcome Results From a Randomized Controlled Trial for</p>	<p>Personality disorder (mixed)</p>	<p>Supportive-Expressive Psychotherapy (SE; Luborsky, 1984)</p>	<p>Non-manualized PDT</p>

	Psychiatric Outpatients with Personality Disorders. <i>Psychotherapy</i> , 46, 362–375. http://www.ncbi.nlm.nih.gov/pubmed/20161588			
183.	Vitriol , V. G., Ballesteros, S. T., Florenzano, R. U., Weil, K. P., & Benadof, D. F. (2009). Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. <i>Psychiatr Serv</i> , 60, 936-942. http://www.ncbi.nlm.nih.gov/pubmed/19564224	Depression (+ childhood trauma)	Brief psychodynamic psychotherapy (Kudler et al, 2003; Davies et al, 1994)	TAU
184.	Watzke , B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., & Schulz, H. (2010). Effectiveness of systematic treatment selection for psychodynamic and cognitive-behavioural therapy: randomised controlled trial in routine mental healthcare. <i>The British Journal of Psychiatry : The Journal of Mental Science</i> , 197(2), 96–105. http://www.ncbi.nlm.nih.gov/pubmed/20679260 Watzke , B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., & Schulz, H. (2012). Longer term outcome of cognitive-behavioural and psychodynamic psychotherapy in routine mental health care: Randomised controlled trial. <i>Behaviour research and therapy</i> , 50(9), 580–587. http://www.ncbi.nlm.nih.gov/pubmed/22750189	Mixed sample	Psychodynamic brief group therapy (Heigl-Evers et al., 1997)	Cognitive-behavioural group therapy (Fiedler, 2001)
185.	Wiborg , I. M., & Dahl, A. A. (1996). Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? <i>Arch Gen Psychiatry</i> , 53, 689-694. http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008	Panic disorder	Brief Dynamic Psychotherapy (BDP; Davanloo, 1978; Malan 1976; 1979; Strupp & Binder, 1984)	Clomipramin alone
186.	Wiltink , J., Dippel, A., Szczepanski, M., Thiede, R., Alt, C., & Beutel, M. E., et al. (2007). Long-term weight loss maintenance after inpatient psychotherapy of severely obese patients based on a randomized study: predictors and maintaining factors of health behavior. <i>J Psychosom Res</i> , 62, 691-698. http://www.ncbi.nlm.nih.gov/pubmed/17540227	Severe obesity	Psychodynamic inpatient treatment (no manual but reference to: Levine & Luborsky, 1981)	Behavioral treatment (no specific reference)
187.	Winston , A., Pollack, J., McCullough, L., Flegenheimer, W., Kestenbaum, R., & Trujillo, M., et al. (1991). Brief psychotherapy of personality disorders. <i>J Nerv Ment Dis</i> , 179, 188-193. http://psycnet.apa.org/psycinfo/1991-25198-001 Winston , A, Laikin, M., Pollack, J., Samstag, L. W., McCullough, L., & Muran, J. C. (1994). Short-term psychotherapy of personality disorders. <i>The American journal of psychiatry</i> , 151(2), 190–4. http://www.ncbi.nlm.nih.gov/pubmed/22217220	Personality disorders (mixed)	Short-Term Dynamic Therapy (STDP; Davanloo, 1980; Also reference to Mann, Malan, and Sifneos)	Brief Adaptive Psychotherapy (BAP; Pollack et al., 1991)
188.	Woody , G. E., Luborsky, L., McLellan, A. T., O'Brien, C. P., Beck, A. T., & Blaine, J., et al.	Opiate dependence	Supportive-Expressive Psychotherapy (SE;	Cognitive-Behavior Therapy (CBT; Beck,

	<p>(1983). Psychotherapy for opiate addicts. Does it help? <i>Arch Gen Psychiatry</i>, 40, 639-645. http://www.ncbi.nlm.nih.gov/pubmed/6847332</p> <p>Woody, G. E., Luborsky, L., McLellan, A. T., & O'Brien, C. P. (1990). Corrections and Revised analyses for Psychotherapy in Methadone Patients. <i>Arch Gen Psychiatry</i>, 47, 788-789. http://www.ncbi.nlm.nih.gov/pubmed/2378551</p>		Luborsky, 1977; unpublished manual)	unpublished) Drug Counseling (DC; Stockdale & Hargrove, unpublished material, 1977)
189.	<p>Woody, G. E., McLellan, A. T., Luborsky, L., & O'Brien, C. P. (1995). Psychotherapy in community methadone programs: A validation study. <i>American Journal of Psychiatry</i>, 152, 1302-1308. http://www.ncbi.nlm.nih.gov/pubmed/7653685</p>	Opiate dependence	Supportive-Expressive Psychotherapy (SE; Luborsky, 1984)	Drug counseling
190.	<p>Zipfel, S., Wild, B., Groß, G., Friederich, H., Teufel, M., Schellberg, D., ... Zwaan, M. De. (2013). Articles Focal psychodynamic therapy, cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. <i>The Lancet</i>, 383, 127-137. http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract</p>	Anorexia	Focal psychodynamic therapy (FPT; Schauenburg et al., 2009)	<p>1. Enhanced cognitive behavior therapy (CBT-E; Fairburn, 2008)</p> <p>2. Optimized TAU</p>